

Chromosomal evidence of species status and evolutionary relationships of the black fly *Prosimulium petrosum* (Diptera, Simuliidae) in Armenia

Sergey Vlasov¹, Maria Harutyunova², Karine Harutyunova², Peter H. Adler³

1 Department of General Biology and Bioecology, Moscow State Regional University, Radio St. 10A, Moscow 105005, Russia **2** Institute of Molecular Biology, Yerevan, Armenia **3** Department of Agricultural and Environmental Sciences, Clemson University, Clemson, SC 29634-0310, U.S.A.

Corresponding author: Sergey Vlasov (sv.vlasov@mgou.ru)

Academic editor: V. Golygina | Received 12 September 2015 | Accepted 24 November 2015 | Published 22 January 2016

<http://zoobank.org/13D1011B-3C3D-4975-8818-DCA81760E8BA>

Citation: Vlasov S, Harutyunova M, Harutyunova K, Adler PH (2016) Chromosomal evidence of species status and evolutionary relationships of the black fly *Prosimulium petrosum* (Diptera, Simuliidae) in Armenia. Comparative Cytogenetics 10(1): 33–44. doi: 10.3897/CompCytogen.v10i1.6551

Abstract

The karyotype of Armenian populations of the black fly *Prosimulium petrosum* Rubtsov, 1955 was characterized and compared with that of all other chromosomally known Palearctic members of the *Prosimulium hirtipes* group. Analysis of the polytene chromosomes established that *Prosimulium petrosum* is most closely related to European populations of *Prosimulium latimucro* (Enderlein, 1925) with which it shares an identical fixed chromosomal banding sequence. Its validity as a species, separate from *Prosimulium latimucro*, is based on its unique sex-differential sections in the expanded centromere region of chromosome I, in agreement with the unique structural configuration of the hypostomal teeth of its larvae. *Prosimulium petrosum* and *Prosimulium latimucro*, therefore, are homosequential species, demonstrating the value of a combined chromosomal and morphological approach in determining species status.

Keywords

Black flies, chromosomal inversions, homosequential species, polytene chromosomes, sex chromosomes

Introduction

Chromosomal rearrangements have long been considered a driving force in speciation in certain groups of organisms, based on a wealth of evidence, much of it indirect (White 1978, Rothfels 1989, Nevo 2012). If the chromosomes have played an integral role in the speciation process, individual species might be expected to carry unique signatures in their karyotype. In the dipteran family Simuliidae, the vast majority of species are chromosomally distinct from one another, even when they cannot be distinguished reliably by morphological criteria (Adler et al. 2010). Detailed banding sequences of the polytene chromosomes in the larval salivary glands of Simuliidae, consequently, have facilitated the discovery of cryptic species, provided insights into population structure and evolutionary relationships, and positioned the Simuliidae at the forefront of knowledge about the genetics of natural populations of insects (Adler and Crosskey 2015a).

The *Prosimulium hirtipes* group is a widespread Holarctic clade of the Simuliidae, consisting of 25 species in the Palearctic Region (Adler and Crosskey 2015b, Adler and Şirin 2015). Twelve of these species have been examined chromosomally, though to various degrees of precision (reviewed by Adler and Crosskey 2015a). Four species of the group occur in Armenia: *Prosimulium frontatum* Terteryan, 1956, *Prosimulium petrosum* Rubtsov, 1955, *Prosimulium rachiliense* Djafarov, 1954, and *Prosimulium tomosvaryi* (Enderlein, 1921) (Adler and Crosskey 2015b). Analyses of the polytene chromosomes of *P. frontatum*, *P. rachiliense*, and *P. tomosvaryi* have revealed cryptic biodiversity and provided hypotheses of their phylogenetic relationships (Adler and Şirin 2014). Comparative chromosomal studies of Armenian populations of *P. petrosum*, however, are lacking, although general karyotypic features, putatively of this species, have been presented for Bulgarian populations (Ralcheva and Dryanovska 1973, Ralcheva 1974, Chubareva and Petrova 2003).

Prosimulium petrosum was described from larvae and pupae collected on 26 May 1952 in Azerbaijan; the holotype larva is from River Agsu above Lake Göygöl (= Geigel) (Rubtsov 1955). Adults attributed to this species were described from Azizbekov (= Vayk) in Armenia (Rubtsov 1956). Terteryan (1968), however, suggested that the descriptions of the Armenian adults represent *P. pronevitshae* Rubtsov, 1955, now a synonym of *P. rachiliense* (Crosskey and Zwick 2007, Adler and Şirin 2014). The pupal gill figured by Rubtsov (1956), based on Azerbaijanian material, has a branching formula of $(2+2+2+2)+(2+2)+(2+2)$, whereas that by Terteryan (1968), based on Armenian material, has a formula of $(3+3+2)+(2+2)+(2+2)$.

Given the lack of chromosomal information for bona fide material of *P. petrosum*, we conducted a comparative band-by-band analysis of *P. petrosum* to characterize its karyotype and illuminate its taxonomic status and evolutionary relationships. In particular, we were interested in determining if *P. petrosum* is a species distinct from the morphologically similar European species, *P. latimucro* (Enderlein, 1925), or if they are conspecific.

Methods

Larvae were collected from three streams, up to about 210 km apart, in April and May in northern and southern Armenia (Table 1). The material was fixed in a 3:1 mixture of ethanol and glacial acetic acid. Pupae and adults were not collected, but 13 mature larvae with well-developed gill histoblasts were obtained. Larvae were identified morphologically as *P. petrosum*, based on structural characters (Rubtsov 1956, Terteryan 1968)—the apex of the median hypostomal tooth of our material was posterior to the apices of the lateral teeth, and the 16 filaments of the pupal gill were arranged on three, widely splayed primary trunks, with a branching formula of (3+3+2)+(2+2)+(2+2) or (3+3+2)+(2+1+1)+(2+2).

Polytene chromosomes from the larval salivary glands were stained using the lacto-aceto orcein method (Bedo 1975), which also stained gonadal tissue. Preparations were spread by squashing chromosomes on a microscope slide. Larval gender was determined by the form of the gonads: rounded in males and elongated in females. Representative chromosomal arms and selected rearrangements were photographed under oil immersion (Figs 1–4). Composite digital images from different focal planes were made with Helicon Focus 5.3 and further processed with Adobe Photoshop CS6. The banding sequences of all six chromosomal arms were compared with the standard maps of the *P. hirtipes* group (Basrur 1962) and with maps of various species in the *P. hirtipes* group (Basrur 1959, Adler and Belqat 2001, Adler and Şirin 2014).

Fixed inversions (i.e., homozygous in all larvae) are italicized in the text and underlined on our maps; floating inversions (i.e., polymorphisms) are not italicized or underlined. Inversions identical to those identified in previous studies (i.e., *IIS-6*, *IIS-7*, *IIL-9*, and *IIL-10*) were given the same numbers assigned by Basrur (1959). Newly discovered inversions were numbered to follow the last number assigned to inversions in other species of the *P. hirtipes* group currently under study and as yet unpublished. Chromosomal terminology, including terms for landmarks, follows that of Basrur (1959, 1962).

Three morphological preparations of mature larvae were deposited in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg, Russia. Additional morphological preparations and chromosomal photographs were deposited in the Moscow State Regional University, Moscow, Russia.

Table 1. Collection data for larvae of *Prosimulium petrosum* in Armenia.

Site	Location	Latitude Longitude	Altitude (m asl)	Date	Larvae analyzed males:females
1	Armenia, Gegarkunik Province, Ddmashen [†]	40°34'N 44°49'E	ca. 1900	21 April 2010	3:5
2	Armenia, Sjunik Province, Mogralzani- Vardanidzor, Megraget River	39°00.40'N 46°12.45'E	ca. 1265	04 May 2011	0:1
3	Armenia, Sjunik Province, Megrinsky pass	39°06.30'N 46°10.47'E	ca. 2375	04 May 2011	13:41

[†] Exact location in Ddmashen area is unknown.

Results

Karyotype. In total, 64 larvae were analyzed. One larva from Site 1 chromosomally matched the banding sequence of *P. rachiliense* cytoform 'A' (*sensu* Adler and Şirin 2014). The other 63 larvae (16 males, 47 females) were assigned to *P. petrosum*. All larvae had a diploid number of $2n = 6$, with tightly paired homologues (Fig. 1).

The chromosomes were submetacentric (Fig. 1). Chromosome I (sections 1–44) was the longest, with the two arms (IS and IL) subequal in length, followed by chromosome II (sections 45–74) with the long arm (IIL) slightly longer than the short arm (IIS). Chromosome III (sections 75–100) was the shortest, with the long arm (IIIL) approximately 35% longer than the short arm (IIIS). The centromere regions of chromosomes I and II were transformed (CI_t , CII_t ; *sensu* Basrur 1959), producing an expanded, flocculent area from section 19 through section 21 (CI_t) and from the middle of section 57 through section 58 (CII_t) (Figs 1–3). The centromere region of chromosome III was not expanded (Figs 1, 4).

A single, primary nucleolus organizer was in the standard position for the *P. hirtipes* group, that is, in the base of IL (Fig. 2). Landmarks that remained in the standard




Figure 1. Total polytene chromosomal complement of *Prosimulium petrosum*. Total polytene chromosomal complement of female larva of *Prosimulium petrosum*, showing the diploid condition of $2n = 6$, with tightly paired homologues.




Figure 2. Chromosome I of *Prosimulium petrosum*, with male and female transformed centromere regions (CI). Breakpoints of autosomal heterozygous inversions are indicated by brackets. C: centromere, NO: nucleolar organizer, 20/21hc: heterochromatic band.

positions for the *P. hirtipes* group included the single Balbiani ring in the base of IIS (Fig. 3), “blister” in IIIS (Fig. 4), and “shield” and “triad” in IIIL (Fig. 4). A chromocenter was lacking, and supernumerary (B) chromosomes were absent.

Fixed (interspecific) inversions. The banding sequence of chromosome arms IS, IL, IIIS, and IIIL was identical with the standard banding sequence established by Basrur (1959, 1962) for the *P. hirtipes* group. Chromosome II, however, had four fixed inversions relative to the standard sequence—two overlapping inversions in the short arm, *IIS-6* and *IIS-7*, and two tandem inversions in the long arm, *IIL-9* and *IIL-10* (Fig. 3). The four inversions involved 73% and 58% of the sections of IIS and IIL, respectively. *IIL-9* moved the “group of 5” marker more centrally and *IIL-10* reversed the polarity of the parabalbani.

Autosomal (intraspecific) polymorphisms. Fifteen autosomal polymorphisms were detected; all were present in the heterozygous state only. These autosomal rear-




Figure 3. Chromosome II of *Prosimulium petrosum*. Relative to the standard sequence, fixed inversions *IIS-6*, *IIS-7*, *IIL-9*, and *IIL-10* are present. Breakpoints of autosomal inversions are indicated by brackets above the chromosomes. BR: Balbiani ring, C: centromere, Pb: parabalbiani, “5”: group of 5 marker.

Table 2. Frequency of homologues with autosomal inversions and other rearrangements (band deletions, duplications, and heterobands) in three Armenian populations of *Prosimulium petrosum*.

Collection site	1	2	3	Armenia [†]
Larvae (<i>n</i>)	8	1	54	63
Chromosomal homologues (<i>n</i>)*	16	2	108	126
IS-27	0.063		0.037	0.040
IS-28	0.063			0.008
IL-16	0.063		0.009	0.016
IIS-16			0.009	0.008
IIL-13			0.009	0.008
IIL-14			0.028	0.024
IIIL-32			0.019	0.016
IIIL-33			0.028	0.024
IIIL-dif [‡]			0.028	0.024
100dlT [‡]			0.046	0.040
90hb			0.019	0.016
87dp			0.009	0.008
Mean number of heterozygous inversions/larva [§]				0.333
Mean number of all heterozygous chromosomal rearrangements/larva [§]				0.460

[†] All three collection sites combined.
^{*} Frequencies of rearrangements were calculated on the basis of the number of homologues.
[‡] Three larvae, which had IIIL-dif (= IIIL-34, IIIL-35+IIIL-36+IIIL-37), also carried heterozygous deletion 100dl_T, the frequency of which is accounted for separately; two additional larvae had heterozygous deletion 100dl_T in the absence of IIIL-dif.
[§] IIIL-dif was treated as a single inversion for the purpose of presenting means.




Figure 4. Chromosome III of *Prosimulium petrosum*. **a, b** chromosome III of *Prosimulium petrosum*. Breakpoints of autosomal inversions and location of 2 additional bands (87dp) are indicated by brackets. IIIIL-dif is an inversion complex, hypothesized to consist of four inversions. Arrows indicate locations of 90hb and 100dlT. **c** heterozygous band duplication 87dp. **d** heteroband 90hb and heterozygous inversion IIIIL-33. **e** complex set of heterozygous inversions, collectively referred to as IIIIL-dif; arrow shows deletion 100dlT in telomere of one homologue. **f** - heterozygous deletion 100dlT; breakpoints of individual inversions IIIIL-34, IIIIL-35+IIIIL-36+IIIIL-37, which comprise complex inversion IIIIL-dif, are indicated by brackets; dashed line designates approximate limits of inversions. C: centromere, bl: blister, ca: capsule, s: shield, tr: triad.

rangements included 12 inversions, one heteroband (90hb), a duplication of bands (87dp), and a telomeric deletion (100dlT) (Table 2, Figs 2, 3, 4). Four inversions in sections 96–100 of IIIIL formed a complex set of loops collectively referred to as IIIIL-dif (Fig. 4, e). The proposed breakpoints of these four inversions are shown in Fig. 4, f. All autosomal polymorphisms were expressed in low frequency (< 0.065; Table 2).

Sex-differential region. All 16 males had a heterochromatic band (20/21hc) at the junction of sections 20 and 21 and lacked conjugation in the CI_t region, typically from section 20 through the beginning of section 21 (Fig. 2), although one male was unpaired from the beginning of section 19 to the beginning of section 21; no inversion could be discerned in the unpaired region. Females lacked the heterochromatic band and exhibited complete pairing of homologues in the CI_t region. Thus, the expanded centromere region of chromosome I was the sex-differential segment, with males X_0Y_1 and females X_0X_0 . In two males, ectopic pairing of CI_t and CII_t occurred in some nuclei.

Discussion

Our chromosomal analysis requires taxonomic context, especially a reasonable assignment of the correct species name. The larvae from our three sites in Armenia are chromosomally cohesive. Based on gill structure, they conform to previous Armenian collections (Terteryan 1968), rather than to Azerbaijani material (Rubtsov 1956), of *P. petrosum*. Based on hypostomal structure, they precisely match the Azerbaijani (holotype) (Rubtsov 1956). Our Armenian collections and the type locality of *P. petrosum* in Azerbaijan are 140–160 km apart, and all are in the same ecoregion—the Caucasus Mixed Forests Ecoregion (World Wildlife Federation 2015). Although a slight difference in the branching pattern of the gill between Armenian and Azerbaijani samples possibly indicates the presence of cryptic species, we attribute the difference to intraspecific variation, which is common, especially on the dorsal trunk, in members of the *P. hirtipes* group (Stloukalova 2004). Similarly, although we found mature larvae 1.0–1.5 months earlier (end of April–beginning of May) than did Terteryan (1968), the seasonal difference could be attributable to altitude or perhaps climatic variation among years. Given the minimal geographic distance, dispersal ability of simuliids (Adler et al. 2005), identical ecoregion, and morphological similarity, we conclude that our Armenian populations are conspecific with the holotype.

Although our material corresponds with the type (Caucasian) concept of *P. petrosum* (Rubtsov 1955), a larger question is whether *P. petrosum* is a unique species or conspecific with *P. latimucro*, as suggested by Adler and Crosskey (2015b), based on morphological similarity. Yankovsky (2003) suggested that the projection of the median hypostomal tooth anterior to the lateral teeth and the second-order branching (i.e., 3+3+2) of the upper gill trunk distinguish the larva of *P. latimucro* from that of *P. petrosum*. Accordingly, our samples correspond with *P. petrosum*, based on the hypostomal teeth, and with *P. latimucro*, based on the branching of the gill.

What do the banding sequences of the polytene chromosomes reveal about possible conspecificity of *P. petrosum* and *P. latimucro* and their evolutionary relationships? The Armenian population of *P. petrosum* shares *IIS-6,7* and *IIL-9* with *P. latimucro*, *P. rufipes* (Meigen, 1830), and *P. “aff. 3”* of Basrur (1959), and fixation of *IIL-10* with *P. rufipes*, *P. “aff. 3”*, and Moroccan *P. latimucro* (Adler and Belqat 2001, Adler and Şirin 2014). In European populations of *P. latimucro*, inversion *IIL-10* is absent or polymorphic. *Prosimulium petrosum* differs from *P. rufipes*, *P. “aff. 3”*, and Moroccan *P. latimucro* by lacking *IS-18*, *IIS-8*, and *IIL-11*, respectively. *Prosimulium petrosum* does not share any autosomal polymorphisms with any studied member of the *P. hirtipes* group. We conclude that European populations of *P. latimucro* are most closely related to *P. petrosum*.

Males and females of *P. petrosum* consistently differ in the expression of their CI_t region, indicating the general location of the sex-determining locus. The sex chromosomes of the Simuliidae often are associated with rearrangements, such as inversions and heterobands, although the X and Y also can be microscopically undifferentiated (X_0Y_0) (Rothfels 1980, Post 1982). Any of the three chromosomes (I, II, or III) can

function as the sex chromosome. Identical, differentiated sex chromosomes are rarely shared between species (Rothfels and Freeman 1983, Adler et al. 2015). Thus, the sex chromosomes can be useful in species discovery and identification (Rothfels 1989).

Lack of pairing of homologues in the CI_t region, observed in males of *P. petrosum*, also is found in at least some populations of other Palearctic members of the *P. hirtipes* group, such as *P. hirtipes* (Fries, 1824), *P. latimucro*, and *P. "aff. 3"*, and often serves as the basis for further elaboration of the Y chromosome, such as the addition of sex-linked inversions and heterobands (Basrur 1959, Adler unpublished). The heterochromatic band 20/21hc of *P. petrosum* also appears on the Y chromosome of various members of the *P. hirtipes* group, including Moroccan populations of *P. rufipes* and *P. latimucro* and some European populations of *P. latimucro* and *P. "aff. 3"*, often with various repatterning of banding in the CI_t region (Adler and Belqat 2001, Adler unpublished). No conspicuous repatterning was observed in the CI_t region of *P. petrosum*. We do not know if the unpaired CI_t condition and heterochromatic band 20/21hc are identical across populations and species, and if so, if their shared nature reflects common ancestry, introgression, or independent origins. Species differences in other members of the *P. hirtipes* group, such as those in eastern North America often are based on minor differences in the centromere region, especially of CIII (Rothfels and Freeman 1977).

A Y chromosome based on an unpaired CI_t region, coupled with 20/21hc, without an associated inversion or band repatterning, uniquely characterizes *P. petrosum*. The allopatric nature of *P. petrosum* and *P. latimucro*, however, presents a challenge for evaluating reproductive isolation; the nearest chromosomally analyzed populations of *P. petrosum* and *P. latimucro* are more than 1,500 km apart. Our analysis of the photographs by Ralcheva (1974) of putative *P. petrosum* from Bulgaria suggests that IS and IL are standard, IIS-6,7 and IIL-9 are present, and IIL-10 is absent; the sex chromosomes and larval morphology were not mentioned. Based on available evidence, *P. petrosum* of Ralcheva (1974), therefore, is probably *P. latimucro*, and most closely resembles populations in the Swiss Alps (as *P. inflatum* "aff. 1" of Basrur 1959). The chromosomal characteristics of all other analyzed populations identified as *P. petrosum* and *P. latimucro* are entirely congruent with the respective configurations of the hypostomal teeth. Thus, we argue that *P. petrosum* is a distinct species on the basis of unique chromosomal features corroborated by distinct hypostomal features. *Prosimulium petrosum* and European *P. latimucro*, therefore, are homosequential species—they have the same fixed banding sequence but differ morphologically, a phenomenon first discovered in *Drosophila* (Carson et al. 1967). Although not common in the Simuliidae, previous examples of homosequential species include several members of the *Simulium vernum* group (Hunter 1987, Adler et al. 2004, Seitz and Adler 2015).

Acknowledgements

This work was supported by the International Scientific and Technical Center (ISTC) Project #A-1662 "Molecular Genetic Monitoring of Blood-Sucking Flies (Diptera)

as a Basis for Biological Control of Vectors of Dangerous Infectious Diseases and Precautions against the Acts of Biological Terrorism”. We thank three reviewers for comments that improved the presentation of the paper.

References

- Adler PH, Belqat B (2001) Cytotaxonomy of the *Prosimulium hirtipes* species group (Diptera: Simuliidae) in Morocco. *Journal of Insect Systematics and Evolution* 32: 411–418. doi: 10.1163/187631201X00281
- Adler PH, Crosskey RW (2015a) Cytotaxonomy of the Simuliidae (Diptera): a systematic and bibliographic conspectus. *Zootaxa* 3975: 1–139. doi: 10.11646/zootaxa.3975.1.1
- Adler PH, Crosskey RW (2015b) World blackflies (Diptera: Simuliidae): a comprehensive revision of the taxonomic and geographical inventory [2015]: 1–123. <http://www.clemson.edu/cafls/biomia/pdfs/blackflyinventory.pdf> [accessed 24 August 2015]
- Adler PH, Şirin Ü (2014) Cytotaxonomy of the *Prosimulium* (Diptera: Simuliidae) of Western Asia. *Zoological Journal of the Linnean Society* 171: 753–768. doi: 10.1111/zoj.12150
- Adler PH, Şirin ÜD (2015) A new species of black fly in the *Prosimulium hirtipes* (Fries) group (Diptera: Simuliidae) from Western Turkey. *Proceedings of the Entomological Society of Washington* 117: 85–94. doi: 10.4289/0013-8797.117.2.85
- Adler PH, Currie DC, Wood DM (2004) The black flies (Simuliidae) of North America. Cornell University Press, Ithaca, NY, xv + 941 pp. + 24 color plates.
- Adler PH, Giberson DJ, Purcell LA (2005) Insular black flies (Diptera: Simuliidae) of North America: tests of colonization hypotheses. *Journal of Biogeography* 32: 211–220. doi: 10.1111/j.1365-2699.2004.01156.x
- Adler PH, Cheke RA, Post RJ (2010) Evolution, epidemiology, and population genetics of black flies (Diptera: Simuliidae). *Infection, Genetics and Evolution* 10: 846–865. doi: 10.1016/j.meegid.2010.07.003
- Adler PH, Cherairia M, Arigue SF, Samraoui B, Belqat B (2015) Cryptic biodiversity in the cytogenome of bird-biting black flies in North Africa. *Medical and Veterinary Entomology* 29: 276–289. doi: 10.1111/mve.12115
- Basrur PK (1959) The salivary gland chromosomes of seven segregates of *Prosimulium* (Diptera: Simuliidae) with a transformed centromere. *Canadian Journal of Zoology* 37: 527–570. doi: 10.1139/z59-059
- Basrur PK (1962) The salivary gland chromosomes of seven species of *Prosimulium* (Diptera: Simuliidae) from Alaska and British Columbia. *Canadian Journal of Zoology* 40: 1019–1033. doi: 10.1139/z62-089
- Bedo DG (1975) C banding in polytene chromosomes of *Simulium ornatipes* and *S. melatum* (Diptera: Simuliidae). *Chromosoma* 51: 291–300. doi: 10.1007/BF00284822
- Carson HL, Clayton FE, Stalker HD (1967) Karyotypic stability and speciation in Hawaiian *Drosophila*. *Proceedings of the National Academy of Sciences USA* 57: 1280–1285. doi: 10.1073/pnas.57.5.1280

- Chubareva LA, Petrova NA (2003) Karyotypes of blackflies (Diptera, Simuliidae) of the world fauna. *Entomologicheskoe Obozrenie* 82: 157–222. [In Russian with English summary; full English translation in *Entomological Review* 83: 149–204.]
- Crosskey RW, Zwick H (2007) New faunal records, with taxonomic annotations, for the blackflies of Turkey (Diptera, Simuliidae). *Aquatic Insects* 29: 21–48. doi: 10.1080/01650420701213853
- Hunter FF (1987) Cytotaxonomy of four European species in the *Eusimulium venum* group (Diptera: Simuliidae). *Canadian Journal of Zoology* 65: 3102–3115. doi: 10.1139/z87-470
- Nevo E (2012) Speciation: chromosomal mechanisms. In: eLS. John Wiley & Sons, Ltd, Chichester. doi: 10.1002/9780470015902.a0001757.pub3
- Post RJ (1982) Sex-linked inversions in blackflies (Diptera: Simuliidae). *Heredity* 48: 85–93. doi: 10.1038/hdy.1982.9
- Ralcheva N[M] (1974) Karyological peculiarities of four species of blackflies of the genus *Prosimulium* Roub., fam. Simuliidae. *Genetika i Selektsiya* [Genetics and Plant Breeding], Sofiya 7: 61–76. [In Bulgarian with English summary]
- Ralcheva NM, Dryanovska OA (1973) Comparative karyological analysis of four species of the genus *Prosimulium* Roub. (Simuliidae). *Doklady Bolgarskoi Akademii Nauk* [Comptes Rendus de l'Académie Bulgare des Sciences] 26: 1085–1088. [In Bulgarian]
- Rothfels KH (1980) Chromosomal variability and speciation in blackflies. In: Blackman RL, Hewitt GM, Ashburner M (Eds) *Insect cytogenetics. Symposium of the Royal Entomological Society of London*. Blackwell Scientific Publications, Oxford 10: 207–224.
- Rothfels K[H] (1989) Speciation in black flies. *Genome* 32: 500–509. doi: 10.1139/g89-475
- Rothfels KH, Freeman DM (1977) The salivary gland chromosomes of seven species of *Prosimulium* (Diptera, Simuliidae) in the *mixture* (*III-1*) group. *Canadian Journal of Zoology* 55: 482–507. doi: 10.1139/z77-065
- Rothfels K[H], Freeman DM (1983) A new species of *Prosimulium* (Diptera: Simuliidae): an interchange as a primary reproductive isolating mechanism? *Canadian Journal of Zoology* 61: 2612–2617. doi: 10.1139/z83-342
- Rubtsov IA (1955) New species of blackflies from Transcaucasia. *Trudy Instituta Zoologii, Akademii Nauk Azerbajjanskoi SSR* 18: 109–129. [In Russian]
- Rubtsov IA (1956) Blackflies (fam. Simuliidae) [Moshki (sem. Simuliidae)]. *Fauna of the USSR. New Series No. 64, Insects, Diptera* 6(6). Akademii Nauk SSSR, Moscow, Leningrad [= Saint Petersburg]. [In Russian; English translation: 1990. Blackflies (Simuliidae). 2nd ed. Fauna of the USSR. Diptera, 6(6). EJ Brill, Leiden, 1042 pp.]
- Seitz G, Adler PH, Forster M (2015) A new species, *Simulium* (*Nevermannia*) *berchtesgadense* (Diptera: Simuliidae), and its chromosomes, from the Alps of southeastern Germany. *Zootaxa* 3937: 248–262. doi: 10.11646/zootaxa.3937.2.2
- Stloukalova V (2004) The variation of pupal gills in *Prosimulium rufipes* (Meigen). *Deutsche Gesellschaft für allgemeine und angewandte Entomologie–Nachrichten* 18: 122–123.
- Terteryan AE (1968) Blackflies (Simuliidae). *Fauna of the Armenian SSR, Diptera* [Moshki (Simuliidae) Fauna Armyanskoi SSR, Nasekomye Dvukrylye]. Izdatel'stvo Akademii Nauk Armyanskoi SSR, Yerevan, Armenia. [In Russian]

White MJD (1978) Modes of speciation. WH Freeman, San Francisco, 455 pp.

World Wildlife Federation (2015) Wildfinder. <http://worldwildlife.org/science/wildfinder/> [accessed 25 August 2015]

Yankovsky AV (2003 [2002]) A key for the identification of blackflies (Diptera: Simuliidae) of Russia and adjacent countries (former USSR). Handbook for the identification of the fauna of Russia 170. Zoologicheskii Institut, Rossiiskaya Akademiya Nauk, Saint Petersburg, 570 pp. [In Russian]